AOM Infusion
Improving Health, Enhancing Life

Gastroenterology Infusion Referral Form

Fax To: 800-528-9860
Phone: 800-746-9089
Website: aominfusionrx.com

Patient Demographic Information

Last Name:		First Name:	
Address:		Apt #:	
City:		State:	Zip:
Primary Phone:		Alt Ph or Email:	
Height:	Weight (kg):	Sex: ☐ Male ☐ Female	DOB:
Emergency Contact:		Phone:	

Insurance Information

Primary Insurance Provider:	Policy Number:
Phone Number:	Group:
Secondary Insurance Provider:	Policy Number:
Phone Number:	Group:
Employer Name:	Phone Number:

Prescription Order

☐ **Remicade or other Infliximab product:** Remicade, Avsola, Inflectra, Renflexis, Unbranded Infliximab	☐ Induction Dose: Infuse: _____mg/kg IV on weeks 0, 2, and 6	☐ Maintenance Dose: Infuse: _____mg/kg IV every 8 weeks ☐ Rapid Infusion: May infuse subsequent doses over one hour based on patient tolerability
☐ **Entyvio**	☐ IV Induction Dose: Infuse Entyvio 300 mg IV over 30 minutes on weeks 0, 2, and 6	☐ Maintenance Dose: Infuse Entyvio 300 mg IV over 30 minutes every 8 weeks ☐ Entyvio Pen Maintenance Dose: After loading dose on week 0 and week 2 of Entyvio 300 mg IV, may switch to 108 mg subcutaneously on week 6 and every 2 weeks thereafter
☐ **Skyrizi**	☐ For Crohn's Disease: Skyrizi 600 mg IV over at least 1 hour at week 0, 4, and 8	☐ For Ulcerative Colitis: Skyrizi 1200 mg IV over at least 2 hours at week 0, 4, and 8
☐ **Stelara or other Ustekinumab product:** Stelara, Steqeyma, Yesintek, Imuldosa, Selarsdi, Pyzchiva, or other biosimilar	☐ Induction Dose: Infuse _____ mg/250 ml NS IV over 1 hour x 1 dose (UC and Crohn's)	☐ Maintenance Dose: Inject _____mg SQ every _____weeks
☐ **Tremfya (IV and Sub Q) for Ulcerative Colitis**	☐ Induction Dose: Infuse Tremfya 200 mg IV over at least one hour with a 0.2 micron filter at week 0, 4, and 8	☐ Maintenance Dose: Inject Tremfya 100 mg PFS SQ at week 16 and every 8 weeks thereafter ☐ Maintenance Dose: Inject Tremfya 200 mg PFS SQ at week 12 and every 4 weeks thereafter. Use the lowest effective recommended dosage to maintain therapeutic response
☐ **Tremfya (IV and Sub Q) for Crohn's Disease**	☐ Induction Dose IV: Infuse Tremfya 200 mg IV over at least one hour with a 0.2 micron filter at week 0, 4, and 8 OR ☐ Induction Dose PFS SQ: 400 mg PFS SQ (as 2 consecutive 200 mg injections) on weeks 0, 4, and 8	☐ Maintenance Dose: Inject Tremfya 100 mg PFS SQ at week 16 and every 8 weeks thereafter ☐ Maintenance Dose: Inject Tremfya 200 mg PFS SQ at week 12 and every 4 weeks thereafter. Use the lowest effective recommended dosage to maintain therapeutic response
☐ **Other Infusion medication (include dosing, frequency, and other pertinent information)** _____		
☐ **Injectafer**	☐ Injectafer 750 mg IV x 2 doses, separated by at least 7 days	
☐ **Venofer**	☐ Venofer 200 mg IV x 5 doses spread over 14 days	

Pre-Medication Order

Pre-Medication: Premedication and anaphylaxis kit per AOM Infusion protocol unless otherwise stated.

☐ Diphenhydramine ☐ PO or ☐ IV _____ mg, approximately 30 minutes prior to each infusion.

☐ Acetaminophen PO_____mg tablet, approximately 30 minutes prior to each infusion.

☐ Methylprednisolone IV_____mg, approximately 30 minutes prior to each infusion.

☐ Other _____ ☐ PO or ☐ IV, Strength _____ Directions _____

☐ Other _____ ☐ PO or ☐ IV, Strength _____ Directions _____

☐ None

Nursing and Lab Orders

Infusion Nursing: RN to provide start of care assessment, place PIV or access CVAD, administer all prescribed medication per Rx, monitor for adverse effects, and teaching until independent with procedure if applicable.

Labs: Lab draw requests will undergo review by the AOM clinical team to assess for current capacity and clinical appropriateness. Please note that we are unable to provide STAT Lab draws or Lab draws on weekends, holidays, or evenings due to specimen stability, logistics, and other considerations.

☐ Lab Order _____ Frequency of Labs: ☐ Once ☐ Every Infusion ☐ Every _____ weeks

☐ Lab Order _____ Frequency of Labs: ☐ Once ☐ Every Infusion ☐ Every _____ weeks

Diagnosis/General Information

Allergies: _____

Pertinent Clinical History: _____

Patient Status (select one):

☐ New to therapy

☐ Continuation of therapy and last infusion day:

_____

Please include the below information in the referral:

☐ Demographics

☐ Insurance Information

☐ Prescription/Order

☐ Clinicals showing infusion need: notes, diagnostics, colonoscopy, medication list, etc

☐ Most Recent TB and Hepatitis B Test Results

☐ Labs

☐ Additional Information: _____

Patient Diagnosis, ICD-10 Codes:

☐ K50.0 Crohn's disease of small intestine

☐ K50.1 Crohn's disease of large intestine

☐ K50.8 Crohn's disease of both small and large intestine

☐ K50.9 Crohn's disease, unspecified

☐ K51.0 Ulcerative (chronic) pancolitis

☐ K51.2 Ulcerative (chronic) proctitis

☐ K51.3 Ulcerative (chronic) rectosigmoiditis

☐ K51.4 Inflammatory Polyps

☐ K51.5 Left sided colitis

☐ K51.8 Other ulcerative colitis

☐ K51.9 Ulcerative colitis, unspecified

☐ Other (code and description):

_____

Prescriber Information

Prescriber:		Office Contact Name:		
Clinic/Practice Name:				
Address:		City:	State:	Zip:
Phone:	Fax:	License #:	UPIN #:	NPI:
Patient Last Name:		Patient First Name:		Patient DOB:

Prescriber Signature: _____ **Date:** _____

Substitutions permitted unless "No Substitution" is noted on this form.

By signing this form, you and your practice are consenting to receive communications from AOM Infusion. You authorize AOM Infusion to assist with prior authorization requests acting as your pharmacy provider in dealing with prescription and medical insurance organizations.

This form is not a valid order in the state of Arizona.

CONFIDENTIALITY NOTICE: The information contained in this communication and any additional attachments is confidential information or Protected Health Information (PHI) belonging to the sender and intended only for the use of the individual or entity named above. If you are not the intended recipient, or the employee or agent responsible for delivering this message to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this communication is strictly prohibited. If you have received this information in error, please notify AOM Infusion immediately at *(800) 556-4246* to arrange for the return or destruction of the original document.

Fax completed form, insurance information, and clinical documentation to: ***800-528-9860***. For any questions, please contact us at ***800-746-9089*** or visit our website ***aominfusionrx.com.***

November 2026 | © 2026 AOM Infusion. All rights reserved.