

### Patient Demographic Information

Last Name:		First Name:	
Address:		Apt #:	
City:	State:	Zip:	
Primary Phone:		Alternate Phone:	
Height:	Weight (kg):	Sex: <input type="checkbox"/> Male <input type="checkbox"/> Female	DOB:
Emergency Contact:		Phone:	

### Insurance Information

Primary Insurance Provider:		Policy Number:	
Phone Number:		Group:	
Secondary Insurance Provider:		Policy Number:	
Phone Number:		Group:	
Employer Name:		Phone Number:	

### Diagnosis/General Information

Primary Diagnosis:		ICD Code:	Caregiver:
Additional Diagnosis:		ICD Code:	Caregiver Phone:
Hx of HTN:	Diabetes:	Allergies:	

### Prescription Information (or attach a copy of the prescription)

**Infusion Therapy:**

Pharmacist will determine appropriate product based on clinical assessment, insurance requirements and availability

OR Preferred Brand \_\_\_\_\_

**Dose:** (please select option(s) and provide complete information, pharmacy to round to the nearest 5 gram vial)

Administration Rate = Follow Manufacturer's Guidelines

Loading Dose: \_\_\_\_\_ gm/kg over \_\_\_\_\_ days, then Maintenance dose: \_\_\_\_\_ gm/kg over \_\_\_\_\_ days, every \_\_\_\_\_ weeks x \_\_\_\_\_ cycles

Infuse: \_\_\_\_\_ gms over \_\_\_\_\_ hours x \_\_\_\_\_ days every \_\_\_\_\_ weeks/months x \_\_\_\_\_ months/cycles

Other Regimen: \_\_\_\_\_

**Infusion Rate:** (please select one and provide complete information)

Pharmacist to determine OR  Start at \_\_\_\_\_ ml/hr, then increase by \_\_\_\_\_ ml/hr every \_\_\_\_\_ minutes to maximum

**Pre-Medication:**

Diphenhydramine, 25 mg capsule: 1-2 capsules by mouth 15-30 minutes before each infusion

Acetaminophen, 325 mg tablet: 1-2 tablets by mouth 15-30 minutes before each infusion

Other \_\_\_\_\_ Strength \_\_\_\_\_ Directions: \_\_\_\_\_

None

**Vascular Access Device:**  Peripheral Catheter  PICC  Port  Other (describe # of lumens) \_\_\_\_\_

**Flush Orders:** (If IV ordered the following flush protocols will be followed):

Sodium Chloride 0.9%

Peripheral Line: 3ml before each dose and 3ml after each dose and prn; Central Line: 5-10ml before each dose and 5-10ml after each dose and prn

Heparin 10 units/ml Peripheral Line: 3ml after last sodium flush and prn

Heparin 100 units/ml Central Line: 5ml after last sodium flush and prn

Provide needles, syringes, VAD supplies & other ancillary supplies needed for infusion

**Hydration Orders:** Infuse \_\_\_\_\_ ml \_\_\_\_\_ solution  Prior to  Following

**Labs:** Results will be faxed to physician's office. If no frequency noted, ordered labs to be done prior to initial infusion only. Labs will not be drawn on weekend/holidays/evenings. Not appropriate for STAT labs.

Quantitative IgA prior to first dispense. Pharmacist to obtain authorization from MD

Other: \_\_\_\_\_  Frequency of Labs: \_\_\_\_\_

**Nursing Orders for Home Infusion Monitor (IV Only)**

**Observe:** Vital signs prior to infusion. Blood pressure and pulse every 15 mins for 1st hour, then every 30 mins until stable infusion rate, then every hour.

**Watch for:** Signs of fluid overload, cardiovascular systems, allergic reactions. **Contact Physician:** For adverse events, stop the infusion. Can restart the infusion at the same or lower rate pending physician's approval or if symptoms subside.

### Physician Information

Prescribing Physician:		Office Contact Name:		
Address:		City:	State:	Zip:
Phone:	Fax:	License #:	UPIN #:	NPI: